



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

Available online at www.sciencedirect.com

ScienceDirect

Biomedical Journal

journal homepage: www.elsevier.com/locate/bj

Letter to the Editor

Imaging abnormalities in pediatric neuro-COVID are more diverse than specified



Josef Finsterer*

Neurology Neurophysiology Center, Vienna, Austria

ARTICLE INFO

Article history:

Received 5 December 2021

Accepted 7 December 2021

Available online 11 December 2021

Keywords:

SARS-CoV-2

COVID-19

Vaccination

GBS

Side effect

Letter to the Editor

We eagerly read the review by Wong and Toh about the neuroimaging findings in pediatric SARS-CoV-2 infections [1]. It was found that pediatric COVID-19 patients can manifest on neuroimaging with encephalitis, AHNE, ADEM, cytotoxic lesion of the corpus callosum, PRES, venous sinus thrombosis, cerebral vasculitis, GBS, transverse myelitis, and myositis [1]. The study is appealing but raises concerns that require discussion.

Several neuro-imaging abnormalities in pediatric COVID-19 patients were not included.

Not addressed in the review was SARS-CoV-2 associated ischemic stroke [2]. SARS-CoV-2 associated ischemic stroke has been reported in a 15yo and 16yo female and being attributed to multisystem inflammatory syndrome in children

(MIS-C), which results from the global immune response to the virus. Both patients responded favourably to immunoglobulins and prednisolone [2].

Not addressed was Wernicke encephalopathy secondary to anosmia and dysgeusia after a SARS-CoV-2 infection [3]. A 15yo female developed intolerance of food intake due to dysgeusia and three months later horizontal gaze palsy, vertical nystagmus, disturbed coordination, and gait ataxia, being attributed to marked thiamine deficiency [3]. FLAIR and DWI modalities of cerebral MRI revealed hyperintensities in the peri-aqueductal grey and dorso-medial thalami, suggesting Wernicke encephalopathy [3].

Not addressed either was pseudotumor cerebri, which has been reported in an 11yo SARS-CoV-2 infected female and manifested with headache, diplopia due to abducens palsy, and papilledema [4].

There was no mentioning of SARS-CoV-2 associated meningitis, which frequently develops in association with encephalitis [5]. or of intra-cerebral haemorrhage and reversible cerebral vaso-constriction syndrome as reported in an 18yo male with COVID-19.

Not discussed was SARS-CoV-2 associated cerebellitis as has been reported in an 11yo male.

Not discussed either was optic neuritis as the presenting manifestation of MIS- C.

Furthermore, there is no reference to subarachnoid bleeding and microbleeds in pediatric COVID-19.

Overall, the interesting review is incomplete and lacks discussion of a number of neuroimaging findings in pediatric COVID-19 patients.

* Corresponding author. Neurology Neurophysiology Center, Postfach 20, 1180 Vienna, Austria.

E-mail address: fifigs1@yahoo.de.

Peer review under responsibility of Chang Gung University.

<https://doi.org/10.1016/j.bj.2021.12.001>

2319-4170/© 2021 Chang Gung University. Publishing services by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Conflicts of interest

The authors declare no conflicts of interest.

REFERENCES

- [1] Wong AMC, Toh CH. Spectrum of neuroimaging mimics in children with COVID-19 infection. *Biomed J* 2022;45:50–62.
- [2] Chang J, Bulwa Z, Breit H, Cherian LJ, Connors JJ, Song SY, et al. Acute large vessel ischemic stroke in patients with COVID-19-related multisystem inflammatory syndrome. *Pediatr Neurol* 2021;126:104–7.
- [3] Landzberg DR, Bery E, Chico S, Koh S, Weissman B. Wernicke encephalopathy from olfactory dysfunction after COVID-19 infection. *Neurologist* 2021;26:274–5.
- [4] Sofuoğlu Aİ, Akçay N, Şevketoğlu E, Bektaş G. Pseudotumor cerebri syndrome as a neurologic involvement of multisystem inflammatory syndrome in children: a case report. *J Trop Pediatr* 2021;67:fmab075.
- [5] Huo L, Xu KL, Wang H. Clinical features of SARS-CoV-2-associated encephalitis and meningitis amid COVID-19 pandemic. *World J Clin Cases* 2021;9:1058–78.